

Three new phragmosporous hyphomycetes on *Ripogonum* from an 'ecological island' in New Zealand

ERIC H.C. MCKENZIE

mckenziee@landcareresearch.co.nz
Manaaki Whenua Landcare Research
Private Bag 92170, Auckland, New Zealand

Abstract — *Corynespora ripogoni* sp. nov., *Ellisembia maungatautari* sp. nov., and *Solicorynespora maungatautari* sp. nov., found on dead stems of *Ripogonum scandens* in New Zealand, are illustrated and described, and compared with related taxa. Two other hyphomycetes are recorded from New Zealand for the first time.

Key words — anamorphic fungi, deuteromycetes, *Ellisembia bambusicola*, *Sporidesmiella parva*, taxonomy

Introduction

Maungatautari Ecological Island is a mainland conservation 'island' in central North Island, New Zealand. It is surrounded by 47 km of pest-proof fence, which was completed in August 2006. Except for a few isolated mice, all mammalian pests within the fence have been eradicated. The first of several threatened species of bird have been re-introduced to the area. During a visit to a small part of the reserve some dead stems of the liana, *Ripogonum scandens* J.R. Forst. & G. Forst. (*Smilacaceae*), were collected. Several phragmosporous species of hyphomycetes were identified from the stems, including three new species that are described below.

Materials and methods

Dead stems of *Ripogonum scandens* were collected from a forested area in the Maungatautari Ecological Island. The stems were incubated under humid conditions and periodically examined for sporulating microfungi. Fungal fruiting structures were removed, mounted in lactophenol or water, and examined by light microscopy. Measurements were made on material mounted in lactophenol. Dried herbarium specimens of the fungi were prepared and deposited in the New Zealand Fungal Herbarium (Herb. PDD). Some other specimens of fungi on *R. scandens* and *R. album* R. Br. (from Australia) held in Herb. PDD were also examined. All of the taxa are treated in alphabetical order by genus.

TABLE 1. Major morphological features of *Corynespora* species described since Siboe et al. (1999)

SPECIES ¹	CONIDIA			Host/ SUBSTRATUM
	Production	Morphology ²	Colour	
<i>acalyphae</i>	Solitary	Obclavate, rostrate	Pale brown to brown	85–120 × 9–11 8–16 <i>Acalypha</i> , Indonesia
<i>albiziicola</i>	Solitary	Obclavate, ellipsoid, or clavate	Pale olivaceous yellow	20–70.1 × 10–18.5 1–6 <i>Albizia</i> , India
<i>aquatica</i>	Solitary	Obclavate to cylindric	Pale brown	34–46 × 3–4.5 (1–)2(–3) Leaves, Mexico
<i>asclepiadacearum</i>	Mostly solitary	Obclavato-cylindric to cylindric	?Pale	44–192 × 10–25 Up to 26 <i>Cryptolepis</i> , India
<i>azadirachtiana</i>	Solitary or catenate	Obclavate	Pale yellow	32–303.5 × 7–21.5 1–20 <i>Azadirachta</i> , India
<i>barleriicola</i>	Solitary	Obclavate to cylindric	Olivaceous yellow	41–246 × 10–18.5 3–14 <i>Barleria</i> , India
<i>beilschmiediae</i>	Solitary	Obclavate	Pale brown to brown	52–144.5 × 8.5–11 7–19 <i>Beilschmiedia</i> , China
<i>bombacearum</i>	Solitary or catenate	Obclavato-cylindric to cylindric	Pale to mid olivaceous	26–206 × 8.5–17 Up to 15 <i>Bombax</i> , India
<i>caryotae</i>	Solitary	Obclavate-elongate	Pinkish brown	45–120 × 6–10 Up to 18 <i>Caryota</i> , Singapore
<i>cassiae</i>	Solitary	Obclavate	Pale brown to olivaceous brown	107.5–214 × 11–14 10–21 <i>Cassia</i> , China
<i>catenulata</i>	Solitary or catenate	Obclavate to obclavato- cylindric	Dark olivaceous yellow to pale olivaceous brown	27.5–225.5 × 11–19 1–24 <i>Clerodendrum</i> , India
<i>colebrookiana</i>	Solitary or catenate	Obclavate, rarely cylindric	Pale yellow	45–330 × 6–22 4–16 <i>Colebrookia</i> , India
<i>cucurbiticola</i>	Solitary or catenate	Obclavato-cylindric	Subhyaline to pale olivaceous	38.5–230 × 6.5–20 6–23 <i>Coccinia</i> , Nepal
<i>curvispora</i>	Solitary or catenate	Narrow obclavate	Straw-coloured to mid brown	40–250 × 10–12 5–10 Herbaceous stems, USA
<i>donacis</i>	Solitary	Obclavate	Olivaceous brown	45–70 × 8–12 10–14 <i>Donax</i> , China
<i>erythropsidis</i>	Solitary	Ellipsoid, doliiform to broad clavate	Pale brown to olivaceous brown	25–31 × 9–12 4 <i>Erythropsis</i> , China
<i>euryae</i>	Solitary	Obclavate	Pale brown to brown	36–67 × 6–9 5–9 <i>Eurya</i> , China
<i>fici-altissimae</i>	Solitary	Obclavate, rostrate	Dark brown	55–85 × 9–12 11–18 <i>Ficus</i> , China
<i>fici-benjaminiae</i>	Solitary	Obclavate	Pale olivaceous brown	51.5–71 × 8–11 5–10 <i>Ficus</i> , China
<i>flagellata</i>	Solitary	Obclavate, rostrate; smooth or verrucose	Dark brown	50–100 × 9–11 5–10 <i>Citrus</i> , Ghana
<i>gorakhpurensis</i>	Solitary	Obclavate to ellipsoid	Pale olivaceous yellow	21–157 × 13–20 3–13 <i>Erythrina</i> , India

<i>gracilis</i>	Solitary	Cylindric to obclavate	Olivaceous	92–138 × 5–7	10–22	<i>Piper</i> , Indonesia
<i>gymnocladi</i>	Solitary	Obclavate	Brown to dark brown	15–40 × 7–10.5	2–6	<i>Gymnocladus</i> , China
<i>hamata</i>	Solitary	Obclavate, hamatate at apex	Pale olivaceous brown	158–198 × 9–11	14–19	Dead wood, Indonesia
<i>holopteleae</i>	Solitary or catenate	Obclavato-cylindric to cylindric	Mid olivaceous	23–234 × 3.6–19.5	0–17	<i>Holoptelea</i> , India
<i>jasminicola</i>	Solitary	Obclavate	Pale olivaceous	39.5–176 × 10–21	2–18	<i>Jasminum</i> , Nepal
<i>kenyensis</i>	Solitary	Obclavate to obpyriform, +/- rostrate	subhyaline to pale brown	60–125 × 16–25	8–15	<i>Sericostachys</i> , Kenya
<i>keskaliicola</i>	Solitary or catenate	Obclavato-cylindric to cylindric	Mid olivaceous	64–164 × 16–28	Up to 17	<i>Hemidesmus</i> , India
<i>laevistipitata</i>	Solitary	Broadly ellipsoid	Red-brown	17.5–24 × 7–8	(0–)1–2 (–3)	<i>Pertusaria</i> (lichen), USA
<i>lasianthi</i>	Solitary	Obclavate, sometimes rostrate	Pale brown to dark brown	50–103.5 × 8.5–10	4–8	<i>Lasianthus</i> , China
<i>leucaenae</i>	Solitary	Obclavate, obovoid or ellipsoid	Pale yellow	16–298 × 10–19	1–28	<i>Leucaena</i> , India
<i>litseae</i>	Solitary	Obclavate	Pale brown to olivaceous brown	105–235 × 10–12	14–34	<i>Litsea</i> , China
<i>merrilliopanacis</i>	Solitary	Obclavate, rostrate	Straw coloured to brown	130–260 × 17–21	12–25	<i>Merrilliopanax</i> , China
<i>micheliae</i>	Solitary	Obclavate, rostrate	Subhyaline to brown	333–360 × 15–19	12–28	<i>Michelia</i> , China
<i>morindae-tinctoriae</i>	Solitary	Obclavate	Pale olivaceous	44–127 × 15–26.5	6–15	<i>Morinda</i> , India
<i>myrioneuronis</i>	Solitary	Obclavate	Pale brown to brown	30–46 × 6.5–8	3–4	<i>Myrioneuron</i> , China
<i>nana</i>	Solitary	Obclavate	Subhyaline to pale olivaceous brown	49.5–110 × 9–18.5	4–14	<i>Lantana</i> , India
<i>parapyrenariae</i>	Solitary	Obclavate	Pale brown to brown	70–100 × 11–14	5–9	<i>Parapyrenaria</i> , China
<i>parvispora</i>	Solitary	Ovoid	Brown	13–15 × 4.5–7.5	1–2	<i>Gynotroches</i> , Singapore
<i>pedaliacearum</i>	Solitary or catenate	Obclavato-cylindric to slightly acicular	Pale olivaceous	16–163 × 3.2–6	3–28	<i>Sesamum</i> , India
<i>phylloshureae</i>	Solitary	Obclavate	Brown	30–50 × 8–10	6–10	<i>Phyllostachys</i> , China
<i>premnigena</i>	Solitary or catenate	Obclavate to obclavato-cylindric	Subhyaline to pale yellow	52–265 × 10–15	1–19	<i>Premna</i> , India
<i>rhapidis-humilis</i>	Solitary	Obclavate, rostrate	Pale to olivaceous brown	90–130 × 6–8	12–16	<i>Rhapis</i> , China
<i>rhododendri</i>	Solitary	Obclavate to long rostrate	Pale brown to olivaceous brown	180–400 × 7.5–11	19–36	<i>Rhododendron</i> , China
<i>ripogoni</i>	Solitary	Obclavate	Brown	60–160 × 10–13.5	7–15	<i>Ripogonum</i> , New Zealand

TABLE 1, concluded.

SPECIES ¹	CONIDIA					HOST/ SUBSTRATUM
	Production	Morphology ²	Colour	Size (µm)	Septation	
<i>rosacearum</i>	Solitary or catenate	Obclavate to obclavato-cylindric	Subhyaline to pale olivaceous	26.5–269 × 9.18.5	1–18	<i>Eriobotrya</i> , India
<i>sacchari</i>	Solitary	Obclavate, rostrate; verrucose or smooth	Pale brown to olivaceous brown	80–120 × 8–9	10–14	<i>Saccharum</i> , China
<i>schleichericola</i>	Solitary or catenate	Obclavate	Pale olivaceous	22.5–66 × 3.8–8.5	1–12	<i>Schleichera</i> , India
<i>scolopiae</i>	Solitary	Obclavate	Pale brown to brown	90–150 × 10–13	8–11	<i>Scolopia</i> , China
<i>sed-acaciae</i>	Solitary	Obclavate	Pale brown to olivaceous brown	40–70 × 11–13.5	8–12	<i>Acacia</i> , China
<i>solani</i>	Solitary or catenate	Obclavate to cylindric	Olivaceous yellow	80.6–276 × 8–10	1–17	<i>Solanum</i> , India
<i>subcylindrica</i>	Catenate	Broadly ellipsoid, subcylindrical	Pale brown	18–60(–90) × 5–13	0–3(–6)	<i>Lippia</i> , Brazil
<i>supkharii</i>	Solitary	Obclavate	Pale olivaceous	22.5–142.5 × 10–17.5	2–11	<i>Phyllanthus</i> , India
<i>tanaceti</i>	Solitary	Obclavate; smooth or verruculose	Pale brown to olivaceous brown	60–104 × 12–16	7–12	<i>Tanacetum</i> , China
<i>tectonae</i>	Solitary	Obclavate, rostrate; verrucose or smooth	Pale brown to olivaceous brown	110–160 × 10–12	12–18	<i>Tectona</i> , China
<i>toonae</i>	Solitary	Obclavate, rostrate	Pale brown to dark brown	65–144 × 7–9	4–14	<i>Toona</i> , China
<i>tremicola</i>	Solitary	Obclavate to ellipsoid	Pale olivaceous	104–296 × 11–16	1–12	<i>Trema</i> , India
<i>trichoides</i>	Solitary	Obclavate-cylindric or obclavate	Pale olivaceous	29–170 × 10–15	3–14	<i>Triumfetta</i> , Nepal
<i>ulmacearum</i>	Solitary	Obclavate	Subhyaline to pale olivaceous	15–106 × 3.5–10	2–16	<i>Trema</i> , India
<i>viticola</i>	Solitary or catenate	Obclavate, cylindric to obovoid	Pale olivaceous	34–170 × 7–17.5	1–14	<i>Cayratia</i> , India
<i>ziziphae</i>	Solitary	Obclavato-cylindric, cylindric, or clavate	Mid olivaceous to straw coloured	33–215 × 10–27	Up to 15	<i>Ziziphus</i> , India

¹ All conidiophores are mononematous and non-stromatic, except for *C. asclepiadacearum* and *C. caroytae*, which are stromatic.

² All conidia are smooth, except where indicated.

Taxonomy

Corynespora

The genus *Corynespora* Güssow is characterised by distoseptate phragmoconidia that are produced through an apical pore in the terminal conidiogenous cell. The conidiogenous cell may proliferate one or more times. The conidia are usually produced singly, but in some species short chains of conidia may form. Siboe et al. (1999) provided a synoptic table of the main morphological features that distinguish 50 accepted species of *Corynespora*. Surprisingly, they omitted their own new species, *C. kenyensis* Siboe et al., from the table. Since then another 61 species have been described in the genus, including a new species, which is described below. To assist with the identification of these additional species, their morphological features are presented in TABLE 1, in a similar format to that used by Siboe et al. (1999). Twenty-three of the new species were described from India or Nepal as the cause of leaf spots on a wide variety of plants (Meenu et al. 1998, Singh et al. 2000, Jain et al. 2002, Sharma et al. 2002a,b, 2003, 2005, Dubey & Rai 2003). From the descriptions and drawings all 23 species appear to be very similar to the common cosmopolitan species, *C. cassicola* (Berk. & M.A. Curtis) C.T. Wei. *Corynespora cassicola* is well known to produce leaf spots on *Carica papaya* and *Cucumis sativus*, and on many other species of plants. A further 22 species have been described from China, principally on dead branches (e.g., Zhang & Xu 2005, Zhang et al. 2009).

Corynespora ripogoni McKenzie, sp. nov.

FIG. 1

MYCOBANK: MB 513214

Coloniae in substrato naturali pilosae, nigrae. Mycelium ex hyphis plerumque in substrato immersum, ramosis, septatis, laevibus, pallide brunneis, tenuitunicatis, 1.5–3.5 µm crassis compositum. Conidiophora macronematosa, mononematosa, erecta, recta vel flexuosa, nonramosa, plerumque 5–7-septata, brunnea, crassitunicata, laevia, 50–115 longa, 6–7 µm crassa, interdum per 1 proliferatione elongascentia. Cellulae conidiogenae monotreticae, in conidiophoris incorporatae, terminales, determinatae, cylindricae, 14–25 µm longa, 4.5–6 µm crassa. Conidia solitaria, sicca, acrogena, brunnea, apicem versus pallidiora, laevia, obclavatae, recta vel leviter flexuosa, 60–180 longa, 10–13.5 µm crassa, apicem versus ad 3–6 µm attenuata, basi truncata 3.5–4.5 µm lata, 7–15-distoseptata, ad septa saepe leniter constricta.

ETYMOLOGY: named after the host substrate, *Ripogonum*.

TYPE: NEW ZEALAND, Waikato, near Pukeatua, Maungatautari Ecological Island, on dead stems of *Ripogonum scandens* (*Smilacaceae*), 12 November 2007, E.H.C. McKenzie (PDD 93526, holotype).

COLONIES on natural substrate hairy, black, consisting of large numbers of individual conidiophores. MYCELIUM mainly immersed in the substratum. HYPHAE branched, septate, smooth, pale brown, thin-walled, 1.5–3.5 µm diam. CONIDIOPHORES differentiated, single, erect, straight or flexuous, unbranched,




FIG. 1. Conidia and conidiogenous cells of *Corynespora ripogoni* (from holotype). Specimens mounted in hydrous lactophenol. Scale bar = 20 μ m.

mainly 5–7-septate, brown, thick-walled, smooth, 50–115 μ m long, 6–7 μ m wide above base, 4.5–6 μ m wide at apex, sometimes with 1 percurrent, enteroblastic apical proliferation. CONIDIOGENOUS CELLS monotretic, integrated, terminal, determinate, cylindrical, 14–25 μ m long, 4.5–6 μ m wide. CONIDIA solitary, dry, acrogenous, brown, paler towards the apex, smooth, obclavate, straight or slightly curved, 60–180 μ m long, 10–13.5 μ m wide in the broadest part (mean = 105.9×11.5 μ m, $n = 25$), 3–6 μ m wide near apex, 3.5–4.5 μ m wide at the protruding truncate base, 7–15-distoseptate, often slightly constricted at some septa (particularly at second septum from base) giving a wavy outline to conidia.

COMMENTS: Several species of *Corynespora* have obclavate conidia that are similar to the conidia of *C. ripogoni* (see Ellis 1971, Siboe et al. 1999, Zhang & Xu 2005). However, *C. ripogoni* can be distinguished from other species by overall shape of the conidia, number of septa, and its conidial dimensions, in particular conidial width. It is morphologically quite distinct from *C. cassicola*.

Ellisembia

The genus *Ellisembia* Subram. is characterised by the formation of solitary, holoblastic conidia on unbranched conidiophores that may undergo percurrent proliferation. The conidia are phragmosporous, often with numerous cells, pale to dark brown or almost black (Subramanian 1992, Wu & Zhuang 2005). Many species of *Ellisembia* were formerly included in *Sporidesmium*. The conidia of *Ellisembia* are distoseptate whereas those of *Sporidesmium* are euseptate (Ellis 1958, 1971, Subramanian 1992, McKenzie 1995). However, *Sporidesmium* and morphologically similar genera, such as *Ellisembia*, are polyphyletic and distributed throughout two major ascomycete classes (Shenoy et al. 2006). The relationships of the New Zealand species are unknown. Robert et al. (2005) list 43 specific names under *Ellisembia*. The species are distinguished primarily on conidial morphology and size (Ellis 1971, Subramanian 1992, Wu & Zhuang 2005). A large-spored specimen collected on *Ripogonum scandens* in New Zealand is distinct from all other known species and is described below. In addition, two other species of *Ellisembia* were found on *R. scandens* in the Maungatautari Ecological Island.

Ellisembia adscendens (Berk.) Subram., Proc. Indian Acad. Sci. B 58 183 (1992).

= *Sporidesmium adscendens* Berk., Ann. Mag. Nat. Hist. 4: 291 (1840).

SPECIMENS EXAMINED: NEW ZEALAND, Auckland, St Heliers Bay, Dingle Dell, on *R. scandens*, 16 April 2004, E.H.C. McKenzie (PDD 80287). Waikato, near Pukeatua, Maungatautari Ecological Island, on *R. scandens*, 12 November 2007, E.H.C. McKenzie (PDD 93263). AUSTRALIA, Queensland, Bunya Mountain National Park, Westcott Plain, on *R. album*, 23 November 1995, E.H.C. McKenzie (PDD 65302).

COMMENTS: This species is widely distributed, usually on woody substrates (Ellis 1971, McKenzie 1995, Wu & Zhuang 2005). In New Zealand it is known only from the northern part of the country on both native and introduced plants.

Ellisembia bambusicola (M.B. Ellis) J. Mena & G. Delgado, in

Mena-Portales et al., Boln Soc. Micol. Madrid 25: 266 (2000).

= *Sporidesmium bambusicola* M.B. Ellis, Mycol. Pap. 70: 34 (1958).

SPECIMEN EXAMINED: NEW ZEALAND, Waikato, near Pukeatua, Maungatautari Ecological Island, on *R. scandens*, 12 November 2007, E.H.C. McKenzie (PDD 94155).

CONIDIOPHORES brown, up to 75 µm long, 7.5 µm wide at base tapering to 4.3 µm wide near apex. CONIDIA brown, 80–110 µm long, 11.5–14 µm wide

at widest point, apex 3.5–5 µm wide, base 4.5–6 µm wide, 10–18-distoseptate, septa averaging 6.3 µm apart.

COMMENTS: The conidial dimensions of the New Zealand specimen are very similar to those given by Ellis (1958) and Wu & Zhuang (2005). Ellis (1958) described the conidia as being 11–25-distoseptate and measuring 65–125 × 11–14 µm, 3–6 µm wide at the apex, and 4–5.5 µm wide at the base. He also stated that the septa averaged 6.2 µm apart. Wu & Zhuang (2005) gave similar measurements, reporting the conidia as 12–20-distoseptate, 60–130 × 13–15 µm, 5–10 µm wide at the apex, and 4–6 µm wide at the base.

This species has not been previously recorded in New Zealand. Superficially the conidia appear similar to those of *E. maungatautari*, especially in overall shape and in the shape of the conico-truncate base. However, the conidia of *E. bambusicola* are considerably smaller and lack the long, hyaline beak. *Ellisembia bambusicola* was originally described on bamboo culms from west Africa. The teleomorph, *Miyoshiella fusispora* Kawam., was described from Japan (Réblová 1999), where it is reported to cause black spots on bamboo. The fungus is also recorded from Cuba, Mexico, USSR, India, Hong Kong, and China (Wu & Zhuang 2005, Farr et al. 2009). Most records are on members of the *Poaceae* and *Areaceae*, although there are reports of the fungus on unidentified branches and trunks (Mena-Portales et al. 2000).

***Ellisembia maungatautari* McKenzie, sp. nov.**

FIG. 2

MYCOBANK: MB 513215

Coloniae in substrato naturali pilosae, nigrae. Mycelium ex hyphis plerumque in substrato immersum. Conidiophora macronematosae, mononematosae, erectae, rectae vel flexuosae, nonramosae, 0–1(–2)-septatae, atro-brunneae, crassitunicatae, laeviae, 15–65 µm longae, 5–9 µm crassae. Cellulae conidiogenerae monoblasticae, in conidiophoris incorporatae, terminales, determinatae, cylindricae, 10–45 µm longae, 6–9 µm crassae. Conidia solitaria, sicca, acrogena, brunnea, plerumque rostrata, laevia vel verruculosis, cylindrica vel obclavata, recta vel leviter flexuosa, 85–125 µm longa (rostrum exclusa), 170–275 µm longa (rostrum inclusa), 13–15 µm crassa, apice acuta, ad basim conico-truncata, 5–7 µm lata, apicem versus ad 0.5–3.5 µm attenuata, 17–23-distoseptata; cellulis basalibus 1 vel 2 majoribus, atro-brunneis; cellulis apicalibus gradatim pallidioribus.

ETYMOLOGY: named after the type locality, Maungatautari Ecological Island.

TYPE: NEW ZEALAND, Waikato, near Pukeatua, Maungatautari Ecological Island, on dead stems of *Ripogonum scandens* (*Smilacaceae*), 12 November 2007, E.H.C. McKenzie (PDD 93259).

COLONIES on natural substrate hairy, black, consisting of large numbers of individual conidiophores. MYCELIUM mainly immersed in the substratum. CONIDIOPHORES differentiated, single, erect, straight or flexuous, unbranched, 0–1(–2)-septate, dark brown, thick-walled, smooth, 15–65 µm long, sometimes of uneven width, 5–6.5 µm wide at apex, 7.5–9 µm wide at mid




FIG. 2. Conidia and conidiogenous cells of *Ellisembia maungatautari* (from holotype). Specimens mounted in hydrous lactophenol. Scale bars = 20 μ m.

point. CONIDIOGENOUS CELLS monoblastic, integrated, terminal, determinate, cylindrical, 10–45 μ m long, 6–9 μ m wide. CONIDIA solitary, dry, acrogenous, brown, 1 or 2 basal cells dark brown, usually rostrate, beak very pale brown, smooth or verruculose, cylindrical or obclavate, straight or slightly curved, 85–125 μ m long (mean = 105 μ m, n = 25) excluding the beak, 170–275 μ m long (mean = 234 μ m, n = 15) including the beak, 13–15 μ m wide in the broadest part (mean = 14.0 μ m, n = 25), 5.5–8 μ m wide at apical septum, 5–7 μ m wide at the conico-truncate base, 17–23-distoseptate; beak 50–80 μ m long, 1.5–3.5 μ m wide at base tapering to 0.5–3.5 μ m wide at apex.

COMMENTS: Several species of *Sporidesmium* sensu lato have conidia with a filiform beak or appendage including *S. magnibrachypus* Matsush., *S. malayasianum* Subram., *S. ochnae* Cheng K. Shi & X.G. Zhang, *S. pruni* Jian Ma & X.G. Zhang, *S. queenslandicum* Matsush., and *S. raphidophorae* K. Zhang & X.G. Zhang. These species are compared with *E. maungatautari* (TABLE 2). *Ellisembia maungatautari* most closely resembles *S. malayasianum* in overall shape, size and colour of the conidia. However, the conidia of *S. malayasianum* are euseptate (Subramanian 1994/95), whereas those of *E. maungatautari* are distoseptate.

TABLE 2. Comparison of *Ellisemia maungatautari* with some morphologically similar species of *Sporidesmium* sensu lato.

SPECIES	CONIDIA				HOST/ SUBSTRATUM
	Septation	Morphology	Size (µm)	Base (µm)	
<i>S. magnibrachypus</i>	9–14 distoseptate	Cylindro-fusiform	48–80* × 12–15.5	3–4 (–5)	Wood, Japan
<i>S. malayasianum</i>	10–17 euseptate	Long subcylindrical to long-subobclavate	160–210 × 10–14	6–9	Palm rachis, Malaysia
<i>E. maungatautari</i>	17–23 distoseptate	Cylindrical to obclavate	170–275 × 13–15	5–7	<i>Ripogonum</i> , New Zealand
<i>S. ochnae</i>	8–11 distoseptate	Obclavate	80–110 × 10–12	4–6	<i>Ochna</i> , China
<i>S. pruni</i>	5–8 distoseptate	Fusiform	45–87 × 8.5–10.5	2–3.6	<i>Prunus</i> , China
<i>S. queenslandicum</i>	10–14 distoseptate	Subulate-cylindrical	60–84 × 7–9	ca 4	<i>Archontophoenix</i> , Australia
<i>S. raphidophorae</i>	14–17 euseptate	Obclavate	93–200 × 12–16.5	4–5.5	<i>Rhaphidophora</i> , China

* excluding beak, which is 60–80 µm long

Helminthosporium palmigenum Matsush., Microfungi Sol. Is. PNG: 30 (1971).

SPECIMENS EXAMINED: NEW ZEALAND, Auckland, Waitakere Ranges, Spragg Bush, on *R. scandens*, November 2007, E.H.C. McKenzie (PDD 54913); Henderson, Corban Estate, banks of Shona Stream, on *R. scandens*, 26 March 2006, E.H.C. McKenzie (PDD 77148).

COMMENTS: This species was first described from Solomon Islands on coconut palm (Matsushima 1971). Since then it has been recorded on several palm genera in Cuba, Venezuela, Taiwan, and Australia (Farr et al. 2009). Surprisingly, in New Zealand *H. palmigenum* is very common on *Ripogonum scandens*; Hughes (1978) cited seven specimens on this host. No specimen was kept from a scant collection from Maungatautari Ecological Island.

Helminthosporium velutinum Link, Ges. Naturf. Freunde Berlin Mag. 3: 10 (1809).

SPECIMENS EXAMINED: NEW ZEALAND, Waikato, near Pukeatua, Maungatautari Ecological Island, on *R. scandens*, 12 November 2007, E.H.C. McKenzie (PDD 93261). Gisborne, Te Urewera National Park, Waikaremoana, Tawa Track, on *R. scandens*, 11 May 2001, E.H.C. McKenzie (PDD 74074, 74080). AUSTRALIA, Queensland, Bunya Mountains National Park, Westcott Plain, on *R. album*, E.H.C. McKenzie (PDD 65299).

COMMENTS: This cosmopolitan species was first recorded from New Zealand by Hughes (1978) on dead wood and bark of several species of native and introduced plants, including *Ripogonum scandens*.

Pseudospiropes simplex (Kunze) M.B. Ellis, Dematiaceous hyphomycetes: 260 (1971).

SPECIMENS EXAMINED: NEW ZEALAND, Auckland, Waitakere Ranges, Spragg Bush, on *R. scandens*, 23 May 1996, E.H.C. McKenzie (PDD 73905). Coromandel, Kauaeranga Valley, Moss Creek Hut Track, on *R. scandens*, 30 August 1986, E.H.C. McKenzie (PDD

52309). Waikato, Mt Pirongia, on *R. scandens*, 21 May 1988, E.H.C. McKenzie (PDD 65789); near Pukeatua, Maungatautari Ecological Island, on *R. scandens*, 12 November 2007, E.H.C. McKenzie (PDD 93266). Wanganui, Kai Iwi, Bushy Park, on *R. scandens*, 15 May 1987, E.H.C. McKenzie (PDD 53614).

COMMENTS: This fungus is common and widespread in New Zealand on wood and bark of a broad range of native and introduced plants (Hughes 1978). It has been recorded previously on *Ripogonum scandens* in New Zealand (Hughes 1978). It is also known from North America, Europe, Africa, and China (Ellis 1971, Shang & Zhang 2007).

Solicorynespora

The genus *Solicorynespora* R.F. Castañeda & W.B. Kendr. was erected to accommodate those *Corynespora*-like species that have euseptate rather than distoseptate conidia (Castañeda Ruíz & Kendrick 1990). While such a character probably does not provide a phylogenetic distinction, it is useful to be able to morphologically divide a relatively large genus such as *Corynespora* into smaller units. Such an approach is followed with *Ellisembia*, which is the distoseptate equivalent of *Sporidesmium*, although the complex has been shown to be polyphyletic (Shenoy et al. 2006). Castañeda Ruíz et al. (2004) provided a key to eight species of *Solicorynespora*, but inexplicably omitted the type species, *S. zapatensis* R.F. Castañeda & W.B. Kendr. and *S. garciniae* (Petch) G. Delgado & J. Mena. Recently, *Corynespora foveolata* (Pat.) S. Hughes was transferred to *Solicorynespora* (Shirouzu & Harada 2008). A new species from Maungatautari Ecological Island is described below.

Solicorynespora maungatautari McKenzie, sp. nov.

FIG. 3

MYCOBANK: MB 513216

Coloniae in substrato naturali effusae, pilosae, nigrae. Mycelium ex hyphis plerumque in substrato immersum, ramosis, septatis, laevibus, luteus vel pallide brunneis, tenuitunicatis, 1.5–2 µm crassis compositum. Conidiophora macronematosa, mononematosa, erecta, recta vel flexuosa, nonramosa, septata, atro-brunnea, apicem versus pallidiora, crassitunicata, laevia, 55–120 µm longa, 3.5–4.5 µm crassa, interdum per 1–2 proliferatione elongascentia. Cellulae conidiogenae monotreticae, in conidiophoris incorporatae, terminales, determinatae, cylindricae, 12–21 µm longa, 3.5–4.5 µm crassa. Conidia solitaria, sicca, acrogena, brunnea, laevia, obclavata vel fusiforma, recta vel leviter flexuosa, 22–41 µm longa, 4.5–6.5 µm crassa, apice acuta, ad basim conico-truncata, 2–2.5 µm lata, apicem versus ad 2–3 µm attenuata, (3–)4–5(–6)-euseptata; cellulis basalibus 2 vel 3 majoribus, brunneis; cellulis apicalibus gradatim pallidioribus.

ETYMOLOGY: named after the type locality, Maungatautari Ecological Island.

TYPE: NEW ZEALAND, Waikato, near Pukeatua, Maungatautari Ecological Island, on dead stems of *Ripogonum scandens* (*Smilacaceae*), 12 November 2007, E.H.C. McKenzie (PDD 93262, holotype).

COLONIES on natural substrate effuse, hairy, black, covering large areas of stem. MYCELIUM mainly immersed in the substratum. HYPHAE branched,



FIG. 3. Conidia and conidiogenous cells of *Solicorynespora maungatautari* (from holotype). Specimens mounted in hydrous lactophenol. Scale bar = 20µm.

septate, smooth, yellowish to pale brown, thin-walled, 1.5–2 µm diam. CONIDIOPHORES differentiated, single, erect, straight or flexuous, unbranched, septate, dark brown, paler towards apex, thick-walled, smooth, 55–120 µm long, 3.5–4.5 µm wide near base, sometimes with 1–2 percurrent, enteroblastic, apical proliferations. CONIDIOGENOUS CELLS monotretic, integrated, terminal, determinate, cylindrical, 12–21 µm long, 3.5–4.5 µm wide. CONIDIA solitary, dry, acrogenous, lower 2 or 3 cells brown, other cells paler, smooth, obclavate to fusiform, gradually tapered to an obtuse apex, more abruptly tapered to a protruding truncate base, straight or slightly curved, 22–41 µm long, 4.5–6.5 µm wide in the broadest part (mean = 30.2×5.9 µm, $n = 25$), 2–2.5 µm wide at the base, 2–3 µm wide near apex, (3–)4–5(–6)-euseptate.

COMMENTS: *Solicorynespora maungatautari* is morphologically most similar to *S. mulanjeensis* (B. Sutton) R.F. Castañeda et al., although conidia of the latter species are much larger ($56\text{--}71 \times 10\text{--}12.5$ µm) with more septa (5–8-euseptate).

Sporidesmiella parva (M.B. Ellis) P.M. Kirk, Trans. Br. Mycol. Soc.

79: 486 (1982) var. *parva*.

≡ *Endophragmia parva* M.B. Ellis, More dematiaceous hyphomycetes: 138 (1976).

SPECIMEN EXAMINED: NEW ZEALAND, Waikato, near Pukeatua, Maungatautari Ecological Island, on *R. scandens*, 12 November 2007, E.H.C. McKenzie (PDD 93260).

CONIDIOPHORES 85–150 μm long, swollen at base up to 9 μm wide, 3.5–4 μm wide above the base tapering to 2.5–3 μm at the apex. CONIDIA 16.5–20.5 $\times$ 3–5.5 μm , 1-distoseptate.

COMMENTS: Ellis (1976) described the conidia of *S. parva* as being 1–2-distoseptate and measuring 15–18 $\times$ 3–4 μm . Kirk (1982) gave slightly broader conidial dimensions of (12–)13–18.5(–20) $\times$ 2.5–4 μm and said the conidia were 1(–2)-distoseptate. In his illustration Kirk drew only one conidium with two septa. When mounted in lactophenol, conidia from the New Zealand specimen appeared to be more than 1-distoseptate; however, in water they were all obviously 1-distoseptate. Wu & Zhuang (2005) recorded conidia from Chinese collections as 16–20 $\times$ 2.5–3 μm and 1–2-distoseptate. *Sporidesmiella parva* has been recorded on a range of host plants from various parts of the world including Cuba, UK, USSR, Malaysia, China, and Japan (Wu & Zhuang 2005). This is the first record of this species from New Zealand.

Acknowledgments

Funds for this research were provided by the New Zealand Foundation for Research, Science and Technology through the Defining New Zealand's Land Biota OBI. Roger Shivas, DPI&F, Indooroopilly, Australia and Steve Stephenson, University of Arkansas, USA are thanked for kindly providing pre-submission peer reviews.

Literature cited

- Castañeda Ruíz RF, Heredia GP, Arias RM, Saikawa M, Minter DW, Stadler M, Guarro J, Decock C. 2004. Two new hyphomycetes from rainforests of México, and *Briansuttonia*, a new genus to accommodate *Corynespora alternarioides*. *Mycotaxon* 89: 297–305.
- Castañeda Ruíz RF, Kendrick WB. 1990. Conidial fungi from Cuba: II. University of Waterloo Biology Series 33: 1–61.
- Dubey RK, Rai AN. 2003. Two new hyphomycetous fungi from India. *Indian Phytopathology* 56: 486–490.
- Ellis MB. 1958. *Clasterosporium* and some allied dematiaceae-phragmosporae. I. *Mycological Papers* 70: 1–89.
- Ellis MB. 1971. Dematiaceous hyphomycetes. Commonwealth Mycological Institute, Kew, U.K. 608 pp.
- Ellis MB. 1976. More dematiaceous hyphomycetes. Commonwealth Mycological Institute, Kew, U.K. 507 pp.
- Farr DF, Rossman AY, Palm ME, McCray EB. 2009. Fungal databases, Systematic Mycology and Microbiology Laboratory, ARS, USDA. Retrieved 12 October 2009, from <http://nt.ars-grin.gov/fungal-databases/>
- Hughes SJ. 1978. New Zealand fungi 25. Miscellaneous species. *New Zealand Journal of Botany* 16: 311–370.
- Jain SL, Rai AN, Mehta P. 2002. Additions to the genus *Corynespora* from India. *Indian Phytopathology* 55: 51–56.
- Kirk PM. 1982. New or interesting microfungi. VI. *Sporidesmiella* gen.nov. (hyphomycetes). *Transactions of the British Mycological Society* 79: 479–489.

- Matsushima T. 1971. Microfungi of the Solomon Islands and Papua-New Guinea. Published by the author, Kobe, Japan.
- McKenzie EHC. 1995. Dematiaceous hyphomycetes on *Pandanaceae*. 5. *Sporidesmium* sensu lato. Mycotaxon 56: 9–29.
- Meenu, Kharwar RN, Bhartiya HD. 1998. Some new forms of genus *Corynespora* from Kathmandu valley of Nepal. Indian Phytopathology 51: 146–151.
- Mena-Portales J, Delgado-Rodríguez G, Heredia-Abarca G. 2000. Nuevas combinaciones para especies de *Sporidesmium* sens. lat. Boletín de la Sociedad Micológica de Madrid 25: 265–269.
- Réblová M. 1999. Studies in *Chaetosphaeria* sensu lato III. *Umbrinosphaeria* gen. nov. and *Miyoshiella* with *Sporidesmium* anamorphs. Mycotaxon 71: 13–43.
- Robert V, Stegehuis G, Stalpers J. 2005. The MycoBank engine and related databases. Retrieved 12 October 2009, from <http://www.mycobank.org>
- Shang Z-Q, Zhang X-G. 2007. Taxonomic studies of *Pseudospiropes* from Yunnan, China. Mycotaxon 100: 149–153.
- Sharma N, Chaudhary S, Kamal. 2002a. Three new species of genus *Corynespora*. Indian Phytopathology 55: 178–181.
- Sharma N, Chaudhary RK, Kamal. 2002b. Five undescribed species of *Corynespora*. Indian Phytopathology 55: 458–463.
- Sharma N, Singh PN, Kamal. 2003. Three new taxa of *Corynespora* causing foliar blight in forest plants of north eastern Uttar Pradesh. Journal of Mycology and Plant Pathology 33: 26–32.
- Sharma N, Soni KK, Jamaluddin, Verma RK. 2005. A new species of *Corynespora* from central India. Indian Phytopathology 58: 503–504.
- Shenoy BD, Jeewon R, Wu WP, Bhat DJ, Hyde KD. 2006. Ribosomal and RPB2 DNA sequence analyses suggest that *Sporidesmium* and morphologically similar genera are polyphyletic. Mycological Research 110: 916–928.
- Shirouzu T, Harada Y. 2008. Lignicolous dematiaceous hyphomycetes in Japan: five new records for Japanese mycoflora, and proposals of a new name, *Helminthosporium magnisporum*, and a new combination, *Solicorynespora foveolata*. Mycoscience 49: 126–131.
- Siboe GM, Kirk PM, Cannon PF. 1999. New dematiaceous hyphomycetes from Kenyan rare plants. Mycotaxon 73: 283–302.
- Singh A, Singh SK, Kamal. 2000. Three new species of *Corynespora* from India. Journal of Mycology and Plant Pathology 30: 44–49.
- Subramanian CV. 1992. A reassessment of *Sporidesmium* (hyphomycetes) and some related taxa. Proceedings of the Indian National Science Academy B 58: 179–190.
- Subramanian CV. 1994/95. Hyphomycetes from south east Asia—novelties from Singapore and Malaysia. Kavaka 22/23: 52–76.
- Wu W, Zhuang W. 2005. *Sporidesmium*, *Endophragmiella* and related genera from China. Fungal Diversity Research Series 15. 351 pp.
- Zhang K, Fu H-B, Zhang X-G. 2009. Taxonomic studies of *Corynespora* from Hainan, China. Mycotaxon 109: 85–93.
- Zhang X-G, Xu J-J. 2005. Taxonomic studies of *Corynespora* from Guangxi, China. Mycotaxon 92: 431–436.